

Key:

- = neutral sample atom or molecule
- ⊕ = positively ionized neutral atom or molecule of sample
- ⊖ = secondary electron associated with ionization event

z = distance from "stop" detector (parallel to the central axis of the drift region)

d = distance ion travels in drift region

V_1 = voltage at "Stop" end of the drift tube

V_2 = voltage at opposite end of drift tube ($V_1 < V_2$)

FIG. 1




FIG. 2

Key:


- = neutral sample atom or molecule
- ⊕ = positively ionized neutral atom or molecule of sample
- ⊙ = secondary electron associated with ionization event

z = distance from "stop" detector (parallel to the central axis of the drift region)

d = distance ion travels in drift region

V_1 = voltage at "Stop" end of the drift tube

V_2 = voltage at opposite end of drift tube ($V_1 < V_2$)






FIG. 4

Key:

- = neutral sample atom or molecule
- ⊕ = positively ionized neutral atom or molecule of sample
- ⊖ = secondary electron associated with ionization event
- z = distance from "stop" detector (parallel to the central axis of the drift region)
- d = distance ion travels in drift region
- V_1 = voltage at "Stop" end of the drift tube
- V_2 = voltage at opposite end of drift tube ($V_1 < V_2$)
- V_3, V_4 = Electrostatic energy analyzer voltages